RECOVERY OF THE RESPIRATORY SYSTEM FOLLOWING BLAST INJURY

Edward G. Damon, John T. Yelverton, Ulrich C. Luft, and Robert K. Jones

THIS WORK SPONSORED BY THE DEFENSE ATOMIC SUPPORT AGENCY UNDER NWER/SUBTASK MA012.

Technical Progress Report

on

Contract No. DA-49-146-XZ-372



THIS WORK, A PHASE OF INVESTIGATIONS DEALING WITH

THE BIOLOGICAL EFFECTS OF BLAST FROM BOMBS, WAS SUPPORTED

BY THE DEFENSE ATOMIC SUPPORT AGENCY OF THE DEPARTMENT OF DEFENSE.

(Reproduction in whole or in part is permitted for any purpose of the United States Government.)

PREPARING AGENCY

Lovelace Foundation for Medical Education and Research

Albuquerque, New Mexico

NATIONAL TECHNICAL INFORMATION SERVICE Springfield, Va. 22151

October 1970

Approved for public release;
Distribution Unlimited

RECOVERY OF THE RESPIRATORY SYSTEM FOLLOWING BLAST INJURY

Edward G. Damon, John T. Yelverton, Ulrich C. Luft, and Robert K. Jones

THIS WORK SPONSORED BY THE DEFENSE ATOMIC SUPPORT AGENCY UNDER NWER/SUBTASK MA012.

Technical Progress Report

on

Contract No. DA-49-146-XZ-372

THE BIOLOGICAL EFFECTS OF BLAST FROM BOMBS, WAS SUPPORTED

BY THE DEFENSE ATOMIC SUPPORT AGENCY OF THE DEPARTMENT OF DEFENSE.

(Reproduction in whole or in part is permitted for any purpose of the United States Government.)

PREPARING AGENCY

Lovelace Foundation for Medical Education and Research
Albuquerque, New Mexico

October 1970

FOREWORD

This report presents the results of studies of the tempolabile response of the pulmonary system of sheep exposed to blast pressures. The study is a sequel to an earlier work in which the acute effects of blast injury on pulmonary function were investigated.

The findings may be of interest to those involved in the treatment of chest injuries, the analysis of weapons effects, or in industrial or military medicine.

This study is a part of a broad program in the field of blast and shock biology designed to obtain data for use in prediction of hazards from explosions and in the development of a sound basis for the prognosis and treatment of blast injuries.

ABSTRACT

The pattern of recovery of the respiratory system from blast injury was investigated in sheep exposed to overpressures in a shock tube. Measurements of the pH and blood gas tensions, determinations of the venous-admixture (Qs/Q) and the alveolar-arterial oxygen gradient (A-a)O2 were conducted before and at intervals up to 132 days following injury. There was an immediate marked increase in Qs/Q, reduction in PaO2, and a moderate increase in (A-a)O2, with very little change in the pH or $P_{\rm CO2}$ of the arterial blood. The greatest recovery was evident within 24 hours with further gradual improvement seen 2, 7, 14, and 21 days after exposure. After the 21st day, most of the animals exhibited virtual complete recovery of the functional efficiency of the pulmonary system as tested at rest.

ACKNOWLEDGMENTS

The authors wish to acknowledge the able technical assistance of Mr. Charles S. Gaylord and Mr. Glen L. Paxton for conducting the shock-tube exposures and obtaining the pressure-time measurements, Dr. Ernest A. Henderson for veterinary surgical support, Mr. Takeshi Minagawa for the illustrations, and Mrs. Berlinda Martinez for typing the manuscript.

Finally, the senior author would like to express sincere appoint of Dr. Donald R. Richmond whose consultations and encouragement greatly facilitated the conduct of this study.

The experimental work discussed in this manuscript was conducted according to the principle enunciated in the "Guide for Laboratory Animal Facilities and Care," prepared by the National Academy of Sciences-National Research Council.

TABLE OF CONTENTS

	-	Page
Forewor	d	i
Abstract		ii
Acknowl	edgments	iii
Introduc	ion	1
	eral	1
Results	• • • • • • • • • • • • • • • • • • • •	2
Discussi	on	2
Referen	es	5
1	Effects of Blast Pressure on Venous-Arterial Shunt $(\mathring{Q}s/\mathring{Q})$ and Arterial Oxygen Tension (Pa_{O_2}) in the Sheep APPENDIX A	3
	TABULATED DATA	
TABLE		
1	Effects of Blast Pressures on Venous-Arterial Saunt in the Sheep	A-2
2	Effects of Blast Pressures on Arterial Oxygen Tension in the Sheep	A-3
3	Effects of Blast Pressures on Arterial Carbon Dioxide Tension in the Sheep	A-4
4	Effects of Blast Pressures on Arterial Blood pH in the Sheep	A-5
5	Effects of Blast Pressures on Alveolar-Arterial Oxygen Difference (A-a)O ₂ in Sheep Breathing Air	8-A

RECOVERY OF THE RESPIRATORY SYSTEM FOLLOWING BLAST INJURY

Edward G. Damon, John T. Yelverton, Ulrich C. Luft, and Robert K. Jones

INTRODUCTION

Most studies of the physiological effects of blast injury to the lungs which have been conducted to date were designed to assess the relation between the extent of lung injury and the immediate or early changes in pulmonary function. 1-8 We are not aware of any investigations in which pulmonary function test parameters have been followed for more than 24 hours following the detonation.

Previous reports have indicated that there is usually an increase in rate and reduction in the depth of respiration occurring soon after exposure. ³
Furthermore, there is an increase in the venous-arterial shunt which generally correlates with the level of blast lung injury. ⁸ The purpose of this study was to extend observations made in the previously cited reference including changes in the pulmonary function in sheep occurring with time after exposure to "sharp"-rising overpressures of "long" duration in the sublethal or near-lethal range.

METHODS

General

Ten young adult ewes having a mean body weight of 40 kg. were exposed singly with their left sides against the endplate of a 42-72-inch-diameter shock tube 5 to "sharp"-rising reflected pressures ranging from 33 to 45 p.s.i.g. with positive-phase durations of 173 to 228 msec. The ambient pressure at exposure was 12 p.s.i.a. 10

Pulmonary function tests, including determinations of the venous-arterial shunt (Qs/Q), alveolar-arterial O_2 gradient (A-a) O_2 , arterial-alveolar (endtidal) CO2 gradient (a-A)CO2, oxygen tension (Pa_{O_2}), carbon dioxide tension (Pa_{CO_2}), and pH of the arterial blood, were conducted without anesthesia from 2 to 13 days prior to exposure, on the day of exposure (usually within 1 hour after the blast), at post-exposure intervals of 1 day, 2, 7, 14, and 21 days and, on some of the animals with patent catheters, at irregular intervals up to 132 days post-shot. All animals were sacrificed for studies of the gross pathology at the end of the study or when the arterial catheter became plugged. The pulmonary-function test procedures have been previously described. Blood samples were drawn from an indwelling arterial catheter inserted through the femoral artery into the aorta. The end-tidal CO2 was measured by a Beckman

Spinco Model LB-1 CO₂ Meter with Linearizer, and the O₂ and CO₂ concentrations of expired gases and calibration gases were determined by the micro Scholander technique. The blood samples were analyzed at 37°C. on an Instrumentation Laboratories Model 113-SI Ultra-Micro pH, P_{O2} , and P_{CO2} Blood Analyzing System, and the values were corrected to the species' body temperature (39°C.).

RESULTS

The major results of the study are given in tables 1 through 5 of appendix A and illustrated in figure 1. The changes in venous-arterial shunt (Qs/Q), expressed as a percent of the cardiac output, are recorded in table 1 of appendix A and illustrated in figure 1(a). The pre-shot mean value was 6.7 with a range of 4.6 to 8.8 percent. Post-shot tests were generally conducted within 1 hour of the exposure, and all animals, except Sheep No. 811, exhibited an increase in Qs/Q at this time. The greatest post-exposure recovery was evident on day 1 with further gradual improvement seen on days 2, 7, and 14. By day 14, four of the eight animals tested had Qs/Q values within the pre-shot range, whereas the other four still had a residual increase in the venous admixture.

A similar pattern of change was evident from the arterial oxygen tension (PaO_2) data listed in table 2 and illustrated in figure 1(b). There was moderate hypoxia immediately after exposure with progressive recovery during the subsequent 14 days.

The data in tables 3 and 4 indicate that there was little or no change in the mean value of the carbon dioxide tension and pH of the arterial blood in these animals. The data on the alveolar-arterial oxygen difference (A-a)O₂, given in table 5, show a mild increase in the mean value of this parameter which reached its peak on the first post-exposure day and returned to pre-exposure levels after the 21st day. There were no consistent changes in the other pulmonary function test parameters recorded during the study.

DISCUSSION

Lung injuries resulting from exposure to blast pressures would be expected to produce pathophysiological effects which, for some undetermined period of time, would become worse due to the progressive development of pulmonary hemorrhage and edema, then stabilize and begin to improve. The improvement in the functional efficiency of the pulmonary system with time may be a twofold process, including an initial physiological compensation involving a reduction in ventilation of non-perfused alveoli and reduction in blood flow through the vasculature of non-ventilated alveoli. The second, and somewhat slower process of recovery, includes the reversal of pulmonary edema, resolution and clearing of pulmonary hemorrhage, and the progressive repair of damaged tissues. The sharp reduction in venous admixture which occurred in these animals 24 hours after exposure may be largely due to compensatory vasoconstriction reducing




Figure 1. --Effects of Blast Pressure on Venous-Arterial Shunt (Qs/Q) and Arterial Oxygen Tension (Pa $_{\rm Q2}$) in the Sheep.

the blood flow through damaged segments of the lung in which the alveolar spaces and airways are filled with hemorrhage and are therefore not open to ventilation; whereas, the slower recovery seen on subsequent days is the result of progressive clearing of hemorrhage, edema, and the restoration of the pulmonary parenchyma through the healing processes. Earlier studies have shown that by 7 days' post-exposure there is marked clearing of the pulmonary hemorrhage. ard by 21 days' post-exposure the lungs are usually entirely clear of hemorrhage. 13 However, one of the sequela of blast injury to the lungs is the occurrence of multiple fibrotic foci, or fine scars, which apparently persist unchanged beyond a period of 2 months. 13 The ultimate fate and functional significance of such scars is unknown. There was very little to indicate a chronic functional effect of the lung injuries sustained by these animals. Although two of the animals exhibited latent increases in the Qs/Q at 94 and 132 days (table 1, Animal Nos. 794 and 771, respectively), the PaO2, PaCO2, pH, and (A-a)O₂ all were within the normal range (tables 2 through 5). It should be emphasized, however, that the animals in this series were exposed to blast pressures in the sublethal or near lettal range and the results might be far different for those surviving exposure to overpressure at or above the midlethal range. Also, the lack of residual effects on the pulmonary function test parameters in this study might be expected because of the tremendous compensatory capacity of the lung, and further resolution of this question must await the use of pulmonary function studies during exercise or other more sensitive tests in the future. Nevertheless, the recovery of the functional efficiency of the pulmonary system within the first 2 to 3 weeks following blast injury is quite remarkable.

REFERENCES

- 1. Hooker, D. R. "Physiological Effects of Air Concussion." Amer. J. Physiol., 67: 219-274, 1924.
- 2. Krohn, P. L., D. Whitteridge, and S. Zuckerman. "Physiological Effects of Blast." Lancet, 1: 252-258, February 28, 1942.
- 3. Denzinger, T. "Physiological Effects of Blast in Air and Water." Chap. XIV-B, German Aviation Medicine, World War II, II: 1225-1259, U.S. Government Printing Office, Washington, D.C., 1950.
- 4. Zuckerman, S. "Experimental Study of Blast Injuries to the Lungs." Lancet, 2: 219-238, 1940.
- 5. Clemedson, C. -J. "An Experimental Study on Air Blast Injuries." Acta Physiol. Scand., 18, Supplement 61: 1-200, 1949.
- 6. Clemedson, C. -J. "Respiration and Pulmonary Gas Exchange in Blast Injury." J. Apply Physiol. 6: 213-220, 1953.
- 7. Clemedson, C. -J., L. Elstorp, H. Pettersson, and A. B. Sundquist. "Changes of Elastic Properties of Lungs of Rabbits in Air Blast Injury." Aerospace Med. 37: 1125-1130, 1966.
- 8. Damon, E. G., J. T. Yelverton, U. C. Luft, K. Mitchell, Jr., and R. K. Jones. "The Acute Effects of Air Blast on Pulmonary Function in Dogs and Sheep." Technical Progress Report No. DASA 2461, Defense Atomic Support Agency, Department of Defense, Washington, D.C., March 1970.
- 9. Richmond, D. R., C. S. Gaylord, and E. G. Damon. "DASA-AEC-Lovelace Foundation Blast-Simulation Facility." Technical Progress Report No. DASA 1853, Defense Atomic Support Agency, Department of Defense, Washington, D.C., August 1966.
- Damon, E. G., C. S. Gaylord, W. Hicks, J. T. Yelverton, D. R. Richmond, and C. S. White. "The Effects of Ambient Pressure on Tolerance of Mammals to Air Blast." Technical Progress Report No. DASA 1852, Defense Atomic Support Agency, Department of Defense, Washington, D.C., August 1966. Also in Aerospace Med. 39: 1039-1047, 1968.
- 11. Yelverton, J. T., E. A. Henderson, and R. W. Dougherty. "A Chronically Implanted Arterial Catheter for Unanesthetized Animals." Cornell Vet. 59: 466-472, July 1969.

- 12. Scholander, P. F. "Analyzer for Accurate Estimation of Respiratory Gases in One-Half Cubic Centimeter Samples." J. Biol. Chem. 167: 235-259, 1947.
- 13. Chiffelle, T. L. "Pathology of Direct Air-Blast Injury." Technical Progress Report No. DASA-1778, Defense Atomic Support Agency, Department of Defense, Washington, D.C., April 1966.

and the second of the second o

The second section of the second section of the second section of the section of

And the second of the second o

on the control of the Market of the Market of the Artificial Control of the Market of the Control of the Contro

and the second s

APPENDIX A TABULATED DATA

TABLE 1

EFFECTS OF BLAST PRESSURES ON VENOUS-ARTERIAL SHUNT IN THE SHEEP

		1000						Ver	, snot	Arteri	Venous Arterial Shunt, Percent	int, P	grcen	44					
				~1 hr						គឺ	Days, Post-Exposure	ost-E	nsodx	7.0					
Xo.	N. C.	Fressure, peig	Shot	Shot	1	2	7	14	2.1	82	35	64	59	99	72	80	98	\$	132
									,										
629	7	*	4.0	7:11		7.6	?	<u>;</u>	2										
821	39	42	5.7	14.6	13.4 12.8		12.5	12.5	12.5 10.9 11.8 10.1	11.8	10.1	8.0	8.7	8.0	8.2	8.	8. 8.		
771	90	40	ñ, Ψ	ۍ و	7.7	7.8		6.5	6.5					:	8.5	-			9.3
750	4	ç	7.3	14.0		14.5	10.1	7.5							37-3-3				
8	7	9	6.7	9.6					· 									_	
¥	×	*	9.9	12.1	10.7	10.4	4.4	10.0	7.8		11.0	9.5						11.3	
1119	£	35	8.5	7.2	5.0												· · · · · · · · · · · · · · · · · · ·		
715	*	*	8.8	15.8	13.4	13.4 13.0 11.4 11.4 10.4	11.4	11.4	10.4		9.0			,	-				
\$2	3	*	7.6	19.3	16.4	13.2	8.6	10.2	7.5		7.3	7.6	9.9		6.1		***************************************		
970	Š	er er	5.1	13.8	6.8	8.8	80.5	3.6	· · · · · · · · · · · · · · · · · · ·										
Mona	0.	36	6.7	12.7	10.4	10.2	10.0	7.0	8.7		9.4	8.3	7.7		9.2				

TABLE 2

EFFECTS OF BLAST PRESSURES ON ARTERIAL OXYGEN TENSION IN THE SHEEP

		Poflector					Arte	rial	Arterial Oxygen Tension (Air), mmHg	n Ten	ion	Air)	E E	Hg					
2.5		Speck		~1 br						Day	s.	Days, Post-Exposure	Kpoeu	re					
Zo.		Pets.	Shot	Shot	1	2	2	7.	12	82	35	49	59	99	72	98	86	*	132
, , ;	7.	,															- 211, 22		
125	7	\$	53	4	2	2	99	20	7.					,					
120	39	42	58	05	51	53	25	53	25		63	8	09	63	55	2	67		
122	ŝ	•	65	47	- 23	56	ii ii	29	65		09	14 14			51				9
750	.		3	53		58	53	63											
8	7	ş	20	5.5					,										
Į,	98	*	9	6.	20	51	25	53	55		*	53						19	:
118	\$	35	. 63	29	62	19		S								****			
715	2	*	+1	46	4	53	7	47	19	• • • •	59	:							
3	•	*	19	7	47	49	79	99	67		19	29	65		3				
970	3	33	23	£	19	79	99	72			+					;			
100	\$	2	3	90	53	55	×	59	82		59	9	63		57) 1	:

TABLE 3

EFFECTS OF BLAST PRESSURES ON ARTERIAL CARBON DIOXIDE TENSION IN THE SHEEP

				<u> </u>			Carr	O uo	Carbon Dioxide Tension (Air), mmHg	Ten	noi e	Air).	mm	Hg	. '	1			
	\$	Shock		~1 pr						Ď	Days, P	Ost-E	Post-Exposure	ure					
Z.	1	Fressure, poig	Strock	Shot	-	2	-	2	12	8	35	49	59	99	72	80	88	2	132
828		45	36	31	32	33	37	32	32										
921	39	42	32	36	31	*	38	33	36		35	39	35	36	33	37	35		
12	80	Ŷ	62	82	27	31		33	<u>*</u>		ጸ				32				33
750	\$	÷	2	33		30	33	9											
8	\$	÷	2	32															
ž	*	2	36	X	37	35	35	33	32		32	33						31	
=	Ç	35	35	2	27	30											:	٠	
715	2	*	31	30	92	33	37	33	33		30								
3	39	*	2	2	62	32	33	32	33		*	32	31		36				-
\$70	Š	8	32	35	33	33	*												<u>. </u>
i į	\$	22	33	33	30	32	35	×	33		33	35	33		2.				

TABLE 4

EFFECTS OF BLAST PRESSURES ON ARTERIAL BLOOD PH IN THE SHEEP

		9.00						A	pH of Arterial Blood (Air)	rteria	11 Blo	V) po	ir)						
		Shock	1	~1 br						Ä	398.	Days, Post-Exposure	Expos	ure					
Mo.	, e	polg.	Shot	Shot	1	2	7	7	21	82	35	49	65	99	72	80	98	2	132
528	:	45	7.47	7.46	7.48	7.42	7.47	7.42 7.47 7.48 7.39	7.39										
23	39	42	7.43	7.43	7.52	7.4	7.45	7.52 7.44 7.45 7.46 7.47	7.47		7.41	7.36	7.46	7.57	7.41 7.36 7.46 7.57 7.43 7.39 7.40	7.39	7.40		
111	•	•	7.45	7.41	7.45 7.43	7.43		7.51 7.45	7.45						7.53				7.41
759	\$	0+	7.47	7.49		7.47	7.47 7.56 7.48	7.48				-	·					,	
8	‡	÷	7.48	7.46		<u></u>													
ĭ	36	*	7.47	7.55	7.47	7.47 7.41	7.41	7.50 7.50	7.50		7.51	7.45						7.45	
911	+ 3	35	7.47	7.45	7.44	7.42													
415	*	*	7.47	7.47	7.29		7.47	7.47 7.50 7.51	7.51		7.42								
3	\$	*	7.48	7.47	7.58	7.52	7.46	7.46 7.47 7.42	7.42		7.42	7.42 7.45 7.40	7.40		7.44				
570	30	33	7.44	7.45	7.41	7.45	7.41 7.45 7.41 7.42	7.42						- - -					
Mean	9	*	7.46	7.46	7.46	7.45	7.46	7.48	7.46	\prod	7.44	7.42	7.43		7.4.7				

TABLE 5

ALVEOLAR-ARTERIAL OXYGEN DIFFERENCE (A-a)O₂ IN SHEEP BREATHING AIR

		Beflected				Iveo	Ar-A	rteri	Alveolar-Arterial Oxygen Difference (Air), mmHg	/gen	Differ	ence	(Air)	mm	H	1			
	Bosy	Shock		#4 1~						Day	Days, P.	Post-Exposure	xpoer	17.6					
No.	N. C.	patg	Shot	Shot	-	2	7	*	12	87	35	64	59	99	72	80	86	ま	132
929	7	45	75	97	42	97	52	\$	×										
621	39	45	33	\$	37	32	45	82	40		7	12	19	*	23	53	18		
1ÎL	9	•	9	32	33	X		23	31		35				39				23
750	7	¢	35	37		35		9	23										
8	7	6	2	3			· · · · · · · · · · · · · · · · · · ·			** <u>, , , , *</u>									
\$	*	2	=	\$	7	32	24	23	39		39	77						23	···
110	\$	35	33	5	52	23							•	,					
118	Я	*		7	\$	7.	37	52	23		33	•							
3	39.	*	7	\$	36	9	2	21	77		5	23	81		2			,	
370	2	66:	•	%	22	*	•	57			····								
Keas	40	36	62	33	35	32	2	*	31		82	91	19		22				

APPENDIX B

DISTRIBUTION

DEPARTMENT OF DEFENSE

Director, Armed Forces Institute of Pathology, Walter Reed Army Medical Center, 14th Street & Alaska Avenue, Washington, D.C. 20012 (1 copy)

Director, Armed Forces Radiobiology Research Institute, Defense Atomic Support Agency, National Naval Medical Center, Bethesda, Maryland 20014 (1 copy)

Assistant to the Secretary of Defense, (Atomic Energy), Washington, D.C. 20301, ATTN: Document Control (1 copy)

Director, Defense Atomic Support Agency, Washington, D.C. 20305,

ATTN: STMD (10 copies)

ATTN: APTL (Technical Library) (2 copies)

ATTN: APSI (Archives) (1 copy)

Administrator, Defense Documentation Center, Cameron Station-Building 5, Alexandria, Virginia 22314, ATTN: Document Control (20 copies)

Director of Defense Research & Engineering, Washington, D.C. 20301, ATTN: Assistant Director (Nuclear Programs), Dr. William W. Carter (1 copy)

Commander, Field Command, Defense Atomic Support Agency, Sandia Base, Albuquerque, New Mexico 87115, ATTN: Technical Library FCWS-SC (1 copy)

Commander, Test Command, Defense Atomic Support Agency, Sandia Base, Albuquerque, New Mexico 87115, ATTN: Document Control (1 copy)

Director, Weapons Systems Evaluation Group, Washington, D. C. 20305, (1 copy)

DEPARTMENT OF THE ARMY

Commandant, Army Medical Field Service School, Brooke Army Medical Center, Fort Sam Houston, Texas 78234, (1 copy)

Director, Army Nuclear Effects Laboratory, Edgewood Arsenal, Maryland 21010, ATTN: Technical Library (1 copy)

Chief, Army Research Office, Life Sciences Division, 3045 Columbia Pike, Arlington, Virginia 22210 (1 copy)

DEPARTMENT OF THE ARMY (Continued)

Director, Ballistic Research Laboratories, Aberdeen Proving Ground, Maryland 21005, ATTN: AMXBR-TB, Mr. Julius J. Meszards (1 copy)

Chief of Research and Development, Department of the Army, Washington, D.C. 20310, ATTN: NCB Division, LTC C. R. Sykes (1 copy)

Director of Civil Defense, Department of the Army, Washington, D.C. 20310, ATTN: Assistant Director for Research (1 copy)

Commanding General, Medical Research and Development Command, Washington, D.C. 20314, ATTN: NBC Sciences Division (1 copy)

Safeguard System Manager, Safeguard System Office, 1320 Wilson Blvd., Arlington, Virginia 22209, ATTN: Plans Division (1 copy)

The Surgeon General, Department of the Army, Washington, D.C. 20360 (1 copy)

Chief, U.S. Army Combat Developments Command, Medical Service Agency, Fort Sam Houston, Texas 78234 (1 copy)

Chief, U.S. Army Combat Developments Command, Chemical-Biological-Radiological Agency, Fort McClellan, Alabama 36205, ATTN: CBR (1 copy)

Commanding Officer, U.S. Army Combat Developments Command, Institute of Nuclear Studies, Fort Bliss, Texas 79916, ATTN: CDINS-E (1 copy)

Director, Walter Reed Army Institute of Research, Walter Reed Army Medical Center, Washington, D.C. 20012, ATTN: Documents Library (1 copy)

DEPARTMENT OF T' 'NAVY

Chief, Bureau of Medicine and Surgery, Navy Department, Washington, D.C. 20390 (1 copy)

Chief of Naval Material, Navy Department, Washington, D.C. 20360,

ATTN: MAT 031 (1 copy)

ATTN: CODE 0312D, I, JAFFE (1 copy)

Chief of Navai Operations, Navy Department, Washington, D.C. 20350, ATTN: OP-752D (1 copy)

Chief of Naval Research, Department of the Navy, Arlington, Virginia 22217,

ATTN: CODE 402 (1 copy)

ATTN: CODE 418, Dr. T. P. Quinn (1 copy)

DEPARTMENT OF THE NAVY (Continued)

Commanding Officer, Naval Aerospace Medical Institute, Naval Aerospace Medical Center, Pensacola, Florida 32512 (1 copy)

Commander, Naval Air Development Center, Johnsville, Warminster, Pennsylvania 18974, ATTN: Code MRPV, Dr. Gloria T. Chisum (1 copy)

Commander, Naval Air Systems Command, Headquarters, Washington, D.C. 20360.

ATTN: AIR 5381 (1 copy) ATTN: AIR 350 (1 copy)

Commanding Officer, Naval Medical Research Institute, National Naval Medical Center, Bethesda, Maryland 20014 (1 copy)

Commander, Naval Ordnance Laboratory, Silver Spring, Maryland 20910,

ATTN: Code 210, Dr. Wayne W. Scanlon (1 copy)
ATTN: Code 213, Mr. Willard L. Derks '1 copy)
ATTN: Code 121, Dr. D. M. Meredith (1 copy)

Director, Naval Research Laboratory, Washington, D.C. 20390, ATTN: Code 2027, Technical Library (1 copy)

Commander, Naval Ship Research and Development Center, Washington, D.C. 20007, ATTN: Library (1 copy)

Commanding Officer, Naval Submarine Medical Center, Naval Submarine Base-New London, Groton, Connecticut 06340 (1 copy)

Commanding Officer, Naval Weapons Evaluation Facility, Kirtland Air Force Base, Albuquerque, New Mexico 87117, ATTN: Document Control (1 copy)

Commander, Naval Weapons Laboratory, Dahlgren, Virginia 22448, ATTN: Mr. W. Culbertson (1 copy)

DEPARTMENT OF THE AIR FORCE

Commander, Aerospace Medical Division, AFSC, Brooks AFB, Texas 78235, ATTN: AMBR (1 copy)

AF Director of Nuclear Safety, Kirtland AFB, New Mexico 87117 (1 copy)

Commander, Air Force Special Weapons Center, AFSC, Kirtland AFB, New Mexico 87117, ATTN: SWTSX, Survivability/Vulnerability Branch (1 copy)

AF Weapons Laboratory, AFSC, Kirtland AFB, New Mexico 87117, ATTN: Chief, WLBR (Biophysics Branch) (1 copy)

DEPARTMENT OF THE AIR FORCE (Continued)

Department of the Air Force, Air Force Office of Scientific Research, 1400 Wilson Blvd., Arlington, Virginia 22209, ATTN: Assistant Executive Director, Research Communications (1 copy)

Commander, Air Force Systems Command, Andrews AFB, Washington, D.C. 20331, ATTN: SGI (1 copy)

Headquarters USAF, Washington, D.C. 20330,

ATTN: AFRDQSN (Strat. & Def. Forces, Nuc. Ord. Div.) (1 copy)

ATTN: AFRDDG (Sci. & Tech. Div., Dir. of Dev.) (1 copy)

USAF Medical Service School, Sheppard AFB, Texas 76311 (1 copy)

USAF Radiological Health Laboratory, AFLC, Wright-Patterson AFB, Ohio 45433 (1 copy)

USAF School of Aerospace Medicine, AFSC, Brooks AFB, Texas 78235,

ATTN: SMRW, Mr. Donald J. Barnes (1 copy)

ATTN: SMRW, Dr. Donald R. Anderson, Chief, Radiobiology Div. (1 copy)

ATTN: SM**OR (1 copy)

6570th Aerospace Medical Research Laboratories, AFSC, Wright-Patterson AFB, Ohio 45433 (1 copy)

6571st Aeromedical Research Laboratory, AFSC, Holloman AFB, New Mexico 88330 (1 copy)

ATOMIC ENERGY COMMISSION

Battelle Memorial Institute, Pacific Northwest Laboratory, P. O. Box 999, Richland, Washington 99352 (1 copy)

Brookhaven National Laboratory, P. O. Box 150, Upton, Long Island, New York 11973, ATTN: Research Library (1 copy)

Los Alamos Scientific Laboratory, P. O. Box 1663, Los Alamos, New Mexico 87544 (1 copy)

Asst. Gen. Mgr. for Military Application, U.S. Atomic Energy Commission, Washington, D.C. 20545, ATTN: Document Control for Research & Development Branch (1 copy)

U.S. Atomic Energy Commission, Washington, D.C. 20545,

ATTN: Document Control for Reports Section (1 copy)

ATTN: Document Control for Bio. & Med. Div., L. J. Deal (150 copies)
ATTN: Document Control for Bio. & Med. Div., H. D. Bruner (1 copy)

ATOMIC ENERGY COMMISSION (Continued)

U.S. Atomic Energy Commission, New York Operations Office, 376 Hudson Street, New York, New York 16014, ATTN: Document Control for Dr. John H. Harley, Director, HASL (1 copy)

U.S. Atomic Energy Commission, Division of Tech. Info. Ext., P. O. Box 62, Oak Ridge, Tennessee 37830, ATTN: Document Control (300 copies)

Union Carbide Corporation, Nuclear Division, Oak Ridge National Laboratory, P. O. Box X, Oak Ridge, Tennessee 37830,

ATTN: Document Control for J. Auxier, Health Physics Dept. (1 copy)
ATTN: Document Control for Atomic & Mol. Processes Info. Ctr. (1 copy)

University of California, Lawrence Radiation Laboratory, Technical Information Division, ν . O. Box 808, Livermore, California 94550, ATTN: Technical Library (Inner Label Only) (1 copy)

OTHER GOVERNMENT

National Aeronautics and Space Administration Rep, Scientific and Technical Info. Facility, P. O. Box 33, College Park, Maryland 20740, ATTN: SAK/DLA-385 (1 copy)

Department of Transportation, National Highway Safety Bureau Research Center, 6th & D Streets, S.W., Washington, D.C. 20591, ATTN: Mr. A. Hirsch (1 copy)

National Library of Medicine, Accessions Branch, 8600 Wisconsin Ave., Bethesda, Maryland 20014 (1 copy)

DEPARTMENT OF DEFENSE CONTRACTORS

Cincinnati General Hospital, Radioisotope Laboratory, Cincinnati, Ohio 45229, ATTN: Dr. Eugene L. Saenger, Director, Radioisotope Laboratory (1 copy)

General Electric Company, Tempo-Center for Advanced Studies, 816 State Street, Santa Barbara, California 93102, ATTN: DASA Information & Analysis Center (1 copy)

Lovelace Foundation for Medical Education and Research, 5200 Gibson Blvd., S.E., Albuquerque, New Mexico 87108,

ATTN: President-Director, Dr. Clayton S. White (1 copy)

ATTN: Assistant Director of Research, Dr. Robert K. Jones (1 copy)

ATTN: Head-Dept of Comp. Environ. Biol., Dr. Donald R. Richmond (1 copy)

ATTN: Associate Scientist, Dr. Edward G. Damon (1 copy) ATTN: Associate Scientist, Dr. E. Royce Fletcher (1 copy)

DEPARTMENT OF DEFENSE CONTRACTORS (Continued)

Medical College of South Carolina, Department of Surgery, 55 Doughty Street, Charleston, South Carolina 29403, ATTN: Associate Professor of Surgery, Dr. Max S. Rittenbury (1 copy)

New York State Veterinary College, Cornell University, Department of Physical Biology, Ithaca, New York 14850, ATTN: Dr. C. L. Comar, Prof./Head, Dept. of Physical Biology (1 copy)

Technology, Inc., Life Sciences Division, 8531 N. New Braunfels Avenue, San Antonio, Texas 78217 (1 copy)

University of California, San Francisco Medical Center, Department of Ophthamology, San Francisco, California 94122, ATTN: Dr. Severin (1 copy)

Virginia Commonwealth University, Medical College of Virginia, Health Science Division, 1200 East Broad Street, Richmond, Virginia 23219, ATTN: Dept. of Biophysics (1 copy)

Recovery of the Respiratory System Following Blast Injury DESCRIPTIVE NOTES (Type of report and inclinates desce) AUTHORIS (Pint mass, middle infinial, instrance) Edward G. Demon, John T. Yelverton, Ulrich C. Luft, and Robert K. Jones REWARD G. Demon, John T. Yelverton, Ulrich C. Luft, and Robert K. Jones REWARD G. Demon, John T. Yelverton, Ulrich C. Luft, and Robert K. Jones T. TOTAL NO. OF PAGES 19 13 13 14. ORIGINATON' REPORT NUMBER(S) A "ROBERT NO. MEER KAOM Task & Subtask AO12 4 Work Unit O1 15. DESTRIBUTION STATEMENT Approved for public release: distribution unlimited. 16. DESTRIBUTION STATEMENT The pattern of recovery of the respiratory system from blast injury was investigated in sheep exposed to overpressures in a shock tube. Regargements of the pH and blood gas tensions, determinations of the venous-admirature (45/4) and the alveolar-arterial oxygen gradient (A-0)c vers-conducted before and at intervals up to 132 days following injury. There was an immediate marked increase in (4/6), reduction in Pago, and a moderate increase in (A-0)c, with very little change in the pH or PCO2 of the arterial blood. The greatest recovery was evident within 24 hours with further gradual improvement seen 2, 7, 14, and 21 days after exposure. After the 21st day, most of the animals exhibited virtual complete recovery of the functional efficiency of the pulmonary system as tested at rest. /	UNCLASSIFIED			
Considerative control of this, body of solution and to be mored when the everel regard to itself the Constantine C				
LARGEDITION (Companies combo) LOUVELAGE POUNDACTION (COMMACTION OF Medical Education & Research Albuquerque, New Mexico LARGEDITION Recovery of the Respiratory System Following Blast Injury LOUVELAGE (Type of report and Section's disco) LOUVELAGE (Type of report and Section's discondition of the Companies of the Section of the Section's discondition of the Companies and Section of the Companies				
LOVELAGE FOUNDATION for Medical Education & Research Albuquerque, New Mexico Albuquerque, New Mexico Approved from the Respiratory System Following Blast Injury Descriptive notes (Type of report and inclusive disce) Loventhus (Print mann, addits indical, last mane) Edward G. Demon, John T. Yelverton, Ulrich C. Luft, and Robert K. Jones REPORT DATE October 1970 DA-19-146-XZ-372 A PROJECT NO. NEER KAOM The Contract of Stanta NO. DA-19-146-XZ-372 A PROJECT NO. NEER KAOM TO CONTRACT ON STANTAN Approved for public release; distribution unlimited. DASA 2580 The pattern of recovery of the respiratory system from blast injury was investing the pattern of recovery of the respiratory system from blast injury was investing sted in sheep exposed to overpressures in a shock tube. Measurgements of the words—arterial oxygen gradient (A-a)C ₂ were conducted before and at intervals up to 132 days following injury. There was an immediate marked increase in \$4/6\$, reduction in Pa _{Q2} , and a moderate increase in (A-a)C ₂ , with very little change in the pH or PC _{Q2} of the arterial blood. The greatest recovery was evident within 24 hours with further gradual improvement seen 2, 7, 14, and 21 days after exposure. After the 21st day, most of the naimals exhibited virtual complete recovery of the functional efficiency of the pulmonary system as tested at rest. /	(Security Classification of Illie, body of abstract ar	nd Indexing ennotation must be		
Recovery of the Respiratory System Following Blast Injury DESCRIPTIVE MOTES (Type of report and inclusive dates) DATECRATE (Type of report and inclusive dates) DATECRATE (Type of report and inclusive dates) DATECRATE (Type of report and inclusive dates) Edward G. Demon, John T. Yelverton, Ulrich C. Luft, and Robert K. Jones DALES (Type of the Respiratory By 19 13 13 13 15 15 19 19 13 13 15 19 19 13 13 15 19 19 19 19 19 19 19 19 19 19 19 19 19		cation & Research		
Recovery of the Respiratory System Following Blast Injury Descriptive mores (Type of separa and inclusive dises) AUTHORIS (Phiet mann, middle initial, lest mann) Edward G. Demon, John T. Yelverton, Ulrich C. Luft, and Robert K. Jones RESPAT DATE OCTOBER 1970 DA-19-116-XZ-372 L. POLICE NO. NIMER KADM Takk & Subtask AD12 WORK Unit O1 DASA 2580 DASA 2580 L. REPORT MOTES (Type of separa middle lest mann) Approved for public release; distribution unlimited. DESTRUCTION STATEMENT Approved for public release; distribution unlimited. DESTRUCTION STATEMENT The pattern of recovery of the respiratory system from blast injury was investigated in sheep exposed to overpressures in a shock tube. Resurgaments of the pH and blood gas tensions, determinations of the venous-admixture (4x/6) and the alveolar-arterial oxygen gradient (A-a)C2 were conducted before and at intervals up to 132 days following injury. There was an immediate marked increase in \$4/6\$, reduction in \$Pag2, and a moderate increase in (A-a)C2, with very little change in the pH or \$PC02. of the arterial blood. The greatest recovery was evident within 24 hours with furthed gradual improvement seen 2, 7, 14, and 21 days after exposure. After the 21st day, most of the unimals exhibited virtual complete recovery of the functional efficiency of the pulmonary system as tested at rest. /	· · · · · · · · · · · · · · · · · · ·			MAGGIF LED
Recovery of the Respiratory System Following Blast Injury DESCRIPTIVE NOTES (Type of report and inclinates desce) AUTHORIS (Pint mass, middle infinial, instrance) Edward G. Demon, John T. Yelverton, Ulrich C. Luft, and Robert K. Jones REWARD G. Demon, John T. Yelverton, Ulrich C. Luft, and Robert K. Jones REWARD G. Demon, John T. Yelverton, Ulrich C. Luft, and Robert K. Jones T. TOTAL NO. OF PAGES 19 13 13 14. ORIGINATON' REPORT NUMBER(S) A "ROBERT NO. MEER KAOM Task & Subtask AO12 4 Work Unit O1 15. DESTRIBUTION STATEMENT Approved for public release: distribution unlimited. 16. DESTRIBUTION STATEMENT The pattern of recovery of the respiratory system from blast injury was investigated in sheep exposed to overpressures in a shock tube. Regargements of the pH and blood gas tensions, determinations of the venous-admirature (45/4) and the alveolar-arterial oxygen gradient (A-0)c vers-conducted before and at intervals up to 132 days following injury. There was an immediate marked increase in (4/6), reduction in Pago, and a moderate increase in (A-0)c, with very little change in the pH or PCO2 of the arterial blood. The greatest recovery was evident within 24 hours with further gradual improvement seen 2, 7, 14, and 21 days after exposure. After the 21st day, most of the animals exhibited virtual complete recovery of the functional efficiency of the pulmonary system as tested at rest. /			1	
Edward G. Demon, John T. Yelverton, Ulrich C. Luft, and Robert K. Jones REFORT DATE October 1970 19 13 **Convert or season no. DA-49-146-XZ-372 **NOUSET NO. NMER KANM **C Task & Subtask A012 **Work Unit 01 **Convert or season no. DA-49-146-XZ-372 **Work Unit 01 **Convert or season no. DA-49-146-XZ-372 **Nouse no. NMER KANM **C Task & Subtask A012 **Work Unit 01 **Convert no. NMER KANM **Approved for public release; distribution unlimited. **It. SUPPLEMENTARY NOTES** **It. SPONSONING MILITARY ACTIVITY Director Defense Atomic Support Agency Washington, D.C. 20305 **The pattern of recovery of the respiratory system from blast injury was investigated in sheep exposed to overpressures in a shock tube. Measurements of the pH and blood gas tensions, determinations of the venous-admixture (4s/4) and the alveolar-arterial oxygen gradient (A-a)C2 were-conducted before and at intervals up to 132 **Approved in sheep exposed to overpressures in a shock tube. Measurements of the pH and blood gas tensions, determinations of the venous-admixture (4s/4) and the alveolar-arterial oxygen gradient (A-a)C2, with very little change in the pH or PCO2 of the arterial blood. The greatest recovery was evident within 24 hours with further gradual improvement seen 2, 7, 14, and 21 days after exposure. After the 21st day, most of the animals exhibited virtual complete recovery of the functional efficiency of the pulmonary system as tested at rest. /	. REPORT TITLE			
Edward G. Demon, John T. Yelverton, Ulrich C. Luft, and Robert K. Jones REFORT DATE October 1970 19 13 **Convert or season no. DA-49-146-XZ-372 **NOUSET NO. NMER KANM **C Task & Subtask A012 **Work Unit 01 **Convert or season no. DA-49-146-XZ-372 **Work Unit 01 **Convert or season no. DA-49-146-XZ-372 **Nouse no. NMER KANM **C Task & Subtask A012 **Work Unit 01 **Convert no. NMER KANM **Approved for public release; distribution unlimited. **It. SUPPLEMENTARY NOTES** **It. SPONSONING MILITARY ACTIVITY Director Defense Atomic Support Agency Washington, D.C. 20305 **The pattern of recovery of the respiratory system from blast injury was investigated in sheep exposed to overpressures in a shock tube. Measurements of the pH and blood gas tensions, determinations of the venous-admixture (4s/4) and the alveolar-arterial oxygen gradient (A-a)C2 were-conducted before and at intervals up to 132 **Approved in sheep exposed to overpressures in a shock tube. Measurements of the pH and blood gas tensions, determinations of the venous-admixture (4s/4) and the alveolar-arterial oxygen gradient (A-a)C2, with very little change in the pH or PCO2 of the arterial blood. The greatest recovery was evident within 24 hours with further gradual improvement seen 2, 7, 14, and 21 days after exposure. After the 21st day, most of the animals exhibited virtual complete recovery of the functional efficiency of the pulmonary system as tested at rest. /	Recovery of the Respiratory System	Following Blast In	dury	
Edward G. Demon, John T. Yelverton, Ulrich C. Luft, and Robert K. Jones REPORT DATE OCTOBER 1970 19 13 13 14 15 16 17 19 13 16 17 18 18 18 18 18 18 18 18 18			UU	(
Edward G. Demon, John T. Yelverton, Ulrich C. Luft, and Robert K. Jones REPORT DATE OCTOBER 1970 19 13 13 14 15 16 17 19 13 16 17 18 18 18 18 18 18 18 18 18				
Edward G. Demon, John T. Yelverton, Ulrich C. Luft, and Robert K. Jones REPORT DATE 72. TOTAL NO. OF PAGES 72. NO. OF REFS 72. NO. OF REFS 72. NO. OF REFS 73. NO. OF REFS 73. NO. OF REFS 74. NO	4. DESCRIPTIVE NOTES (Type of seport and inclusive dates	•)		
October 1970 DA-49-146-7Z-372 A PROJECT NO. MYER XAXM Task & Subtask A012 Approved for public release; distribution unlimited. DASA 2580 DECENTRACY The pattern of recovery of the respiratory system from blast injury was investigated in sheep exposed to overpressures in a shock tube. Measurements of the pH and blood gas tensions, determinations of the vencus-admirture (45/4) and the alveolar-arterial oxygen gradient (A-a)C ₂ were-conducted before and at intervals up to 132 days following injury. There was an immediate marked increase in \$a/4\$, reduction in Pa _{O2} , and a moderate increase in (A-a)C ₂ , with very little change in the pH or P _{O2} of the arterial blood. The greatest recovery was evident within 24 hours with furthe gradual improvement seem 2, 7, 14, and 21 days after exposure. After the 21st day, most of the animals exhibited virtual complete recovery of the functional efficiency of the pulmonary system as tested at rest.	S- AUTHOR(S) (First name, middle initial, last name)			
October 1970 DA-49-146-7Z-372 A PROJECT NO. MYER XAXM Task & Subtask A012 Approved for public release; distribution unlimited. DASA 2580 DECENTRACY The pattern of recovery of the respiratory system from blast injury was investigated in sheep exposed to overpressures in a shock tube. Measurements of the pH and blood gas tensions, determinations of the vencus-admirture (45/4) and the alveolar-arterial oxygen gradient (A-a)C ₂ were-conducted before and at intervals up to 132 days following injury. There was an immediate marked increase in \$a/4\$, reduction in Pa _{O2} , and a moderate increase in (A-a)C ₂ , with very little change in the pH or P _{O2} of the arterial blood. The greatest recovery was evident within 24 hours with furthe gradual improvement seem 2, 7, 14, and 21 days after exposure. After the 21st day, most of the animals exhibited virtual complete recovery of the functional efficiency of the pulmonary system as tested at rest.	Edward G. Demon. John T. Yelverton	. Ulrich C. Luft. s	nd Robert	K. Jones
October 1970 DA-M-9-IM-CXZ-372 A PROPECT NO. NWER XAXM Task & Subtask A012 A Work Unit 01 Approved for public release; distribution unlimited. DASA 2580 DASA 2580 11. SUPPLEMENTARY NOTES The pattern of recovery of the respiratory system from blast injury was investigated in sheep exposed to overpressures in a shock tube. Measurements of the pR and blood gas tensions, determinations of the vencus-admixture (40/4) and the alveolar-arterial oxygen gradient (A-a)02 were-tonducted before and at intervals up to 132 days following injury. There was an immediate marked increase in Qs/4, reduction in Pao2, and a moderate increase in (A-a)02, with very little change in the pB or POO2 of the arterial blood. The greatest recovery was evident within 24 hours with furthe gradual improvement seen 2, 7, 14, and 21 days after exposure. After the 21st day, of the pulmonary system as tested at rest.	, , , , , , , , , , , , , , , , , , , ,	,, .		
October 1970 DA-M-9-IM-CXZ-372 A PROPECT NO. NWER XAXM Task & Subtask A012 A Work Unit 01 Approved for public release; distribution unlimited. DASA 2580 DASA 2580 11. SUPPLEMENTARY NOTES The pattern of recovery of the respiratory system from blast injury was investigated in sheep exposed to overpressures in a shock tube. Measurements of the pR and blood gas tensions, determinations of the vencus-admixture (40/4) and the alveolar-arterial oxygen gradient (A-a)02 were-tonducted before and at intervals up to 132 days following injury. There was an immediate marked increase in Qs/4, reduction in Pao2, and a moderate increase in (A-a)02, with very little change in the pB or POO2 of the arterial blood. The greatest recovery was evident within 24 hours with furthe gradual improvement seen 2, 7, 14, and 21 days after exposure. After the 21st day, of the pulmonary system as tested at rest.				
DA-49-146-XZ-372 DA-49-146-XZ-372 A "POWER NAME AND DASA 2580 The pattern of recovery of the respiratory system from blast injury was investigated in sheep exposed to overpressures in a shock tube. Heapurgments of the pH and blood gas tensions, determinations of the venous-admixture (JP/Q) and a moderate increase in (A-a)O ₂ , with very little change in the pH or POO ₂ of the arterial blood. The greatest recovery was evident within 24 hours with further gradual improvement seen 2, 7, 14, and 21 days after exposure. After the 21st day, most of the pulmonary system as tested at rest. DEMARKSTEED TARGETTED	. REPORT DATE	78. TOTAL NO. O	PAGES	75. NO. OF REFS
DA-49-146-XZ-372 A POLICY NO. NMER XAXM C. Task & Subtask A012 A Work Unit 01 DASA 2580 DASA 2580 DASA 2580 DASA 2580 DEFINISHED THE STATEMENT Approved for public release; distribution unlimited. The pattern of recovery of the respiratory system from blast injury was investigated in sheep exposed to overpressures in a shock tube. Measurements of the pH and blood gas tensions, determinations of the vencus-admixture (4s/Q) and the alveolar-arterial oxygen gradient (A-a)C2 were-conducted before and at intervals up to 132 days following injury. There was an immediate marked increase in (s/Q, reduction in PaQ2, and a moderate increase in (A-a)C2, with very little change in the pH or PCO2 of the arterial blood. The greatest recovery was evident within 24 hours with further the 21st days, most of the enimals exhibited virtual complete recovery of the functional efficiency of the pulmonary system as tested at rest.	_ ·			<u></u>
a. Task & Subtask A012 d. Work Unit Ol Approved for public release; distribution unlimited. The pattern of recovery of the respiratory system from blast injury was investigated in sheep exposed to overpressures in a shock tube. Measurgeents of the pH and blood gas tensions, determinations of the vencus-admixture (45/Q) and the alveolar-arterial oxygen gradient (A-a)C ₂ were-conducted before and at intervals up to 132 days following injury. There was an immediate marked increase in Qs/Q, reduction in Pa _{O2} , and a moderate increase in (A-a)C ₂ , with very little change in the pH or PCO ₂ of the arterial blood. The greatest recovery was evident within 24 hours with further gradual improvement seen 2, 7, 14, and 21 days after exposure. After the 21st day, most of the animals exhibited virtual complete recovery of the functional efficiency of the pulmonary system as tested at rest.		Se. ORIGINATOR	REPORT NUM	BER(S)
a Work Unit 01 Approved for public release; distribution unlimited. Laborator The pattern of recovery of the respiratory system from blast injury was investigated in sheep exposed to overpressures in a shock tube. Negargements of the pH and blood gas tensions, determinations of the vencus-admixture (as/Q) and the alveolar-arterial oxygen gradient (A-a)02 were conducted before and at intervals up to 132 days following injury. There was an immediate marked increase in Qs/Q, reduction in Pao2, and a moderate increase in (A-a)02, with very little change in the pH or Pco2 of the arterial blood. The greatest recovery was evident within 24 hours with further gradual improvement seem 2, 7, 14, and 21 days after exposure. After the 21st day, most of the animals exhibited virtual complete recovery of the functional efficiency of the pulmonary system as tested at rest.		İ		
Approved for public release; distribution unlimited. Approved for public release; distribution unlimited. It supplementant notice It supplements of the percent notice It supplements notice It supplement	S. PROJECT NO. NWER XAXM			
Approved for public release; distribution unlimited. Approved for public release; distribution unlimited. It supplementant notice It supplements of the percent notice It supplements notice It supplement	. Task & Subtask A012	A OTHER RES	BT NO(S) (Ann	the numbers that was he seed mad
Approved for public release; distribution unlimited. Complementary notes	a first a strading Nows	this report)		value interior and any be accepted
Approved for public release; distribution unlimited. 12. SPONSORING MILITARY ACTIVITY Director Defense Atomic Support Agency Washington, D.C. 20305 Assistance The pattern of recovery of the respiratory system from blast injury was investigated in sheep exposed to overpressures in a shock tube. Measurgments of the pH and blood gas tensions, determinations of the vencus-admixture (49/Q) and the alveolar-arterial oxygen gradient (A-a)02 were conducted before and at intervals up to 132 days following injury. There was an immediate marked increase in 49/Q, reduction in Pao2, and a moderate increase in (A-a)02, with very little change in the pH or Pco2 of the arterial blood. The greatest recovery was evident within 24 hours with furthe gradual improvement seen 2, 7, 14, and 21 days after exposure. After the 21st day, most of the animals exhibited virtual complete recovery of the functional efficiency of the pulmonary system as tested at rest. /	d Work Unit Ol	DA DA	SA 2580	
Director The pattern of recovery of the respiratory system from blast injury was investigated in sheep exposed to overpressures in a shock tube. Measurements of the pH and blood gas tensions, determinations of the vencus-admixture (45/4) and the alveolaraterial oxygen gradient (A-a)C ₂ were-conducted before and at intervals up to 132 days following injury. There was an immediate marked increase in \(\frac{1}{2} \), reduction in Pa _{O2} , and a moderate increase in \((A-a)C_2 \), with very little change in the pH or P _{OO2} of the arterial blood. The greatest recovery was evident within 24 hours with furthe gradual improvement seen 2, 7, 14, and 21 days after exposure. After the 21st day, most of the animals exhibited virtual complete recovery of the functional efficiency of the pulmonary system as tested at rest. /	10. DISTRIBUTION STATEMENT			
Director The pattern of recovery of the respiratory system from blast injury was investigated in sheep exposed to overpressures in a shock tube. Measurements of the pH and blood gas tensions, determinations of the vencus-admixture (45/4) and the alveolaraterial oxygen gradient (A-a)C ₂ were-conducted before and at intervals up to 132 days following injury. There was an immediate marked increase in \(\frac{1}{2} \), reduction in Pa _{O2} , and a moderate increase in \((A-a)C_2 \), with very little change in the pH or P _{OO2} of the arterial blood. The greatest recovery was evident within 24 hours with furthe gradual improvement seen 2, 7, 14, and 21 days after exposure. After the 21st day, most of the animals exhibited virtual complete recovery of the functional efficiency of the pulmonary system as tested at rest. /				
Director Defense Atomic Support Agency Washington, D.C. 20305 The pattern of recovery of the respiratory system from blast injury was investigated in sheep exposed to overpressures in a shock tube. Measurements of the pH and blood gas tensions, determinations of the vencus-admixture (45/4) and the alveolar-arterial oxygen gradient (A-a)C ₂ were conducted before and at intervals up to 132 days following injury. There was an immediate marked increase in Qs/4, reduction in Pa _{O2} , and a moderate increase in (A-a)C ₂ , with very little change in the pH or P _{CO2} of the arterial blood. The greatest recovery was evident within 24 hours with furthe gradual improvement seen 2, 7, 14, and 21 days after exposure. After the 21st day, most of the animals exhibited virtual complete recovery of the functional efficiency of the pulmonary system as tested at rest.				
Director Defense Atomic Support Agency Washington, D.C. 20305 The pattern of recovery of the respiratory system from blast injury was investigated in sheep exposed to overpressures in a shock tube. Measurements of the pH and blood gas tensions, determinations of the vencus-admixture (4/2/Q) and the alveolar-arterial oxygen gradient (A-a)02 were conducted before and at intervals up to 132 days following injury. There was an immediate marked increase in Qs/Q, reduction in Pa _{Q2} , and a moderate increase in (A-a)02, with very little change in the pH or P _{QQ2} of the arterial blood. The greatest recovery was evident within 24 hours with further gradual improvement seen 2, 7, 14, and 21 days after exposure. After the 21st day, most of the animals exhibited virtual complete recovery of the functional efficiency of the pulmonary system as tested at rest.	distribution uni	limited.		
Defense Atomic Support Agency Washington, D.C. 20305 The pattern of recovery of the respiratory system from blast injury was investigated in sheep exposed to overpressures in a shock tube. Measurements of the pH and blood gas tensions, determinations of the vencus-admixture (45/4) and the alveolar-arterial oxygen gradient (A-a)C ₂ were conducted before and at intervals up to 132 days following injury. There was an immediate marked increase in Qs/4, reduction in Pa _{O2} , and a moderate increase in (A-a)O ₂ , with very little change in the pH or PcO ₂ of the arterial blood. The greatest recovery was evident within 24 hours with furthe gradual improvement seem 2, 7, 14, and 21 days after exposure. After the 21st day, most of the animals exhibited virtual complete recovery of the functional efficiency of the pulmonary system as tested at rest.	II. SUPPLEMENTARY NOTES		MILITARY ACT	IVITY
The pattern of recovery of the respiratory system from blast injury was investigated in sheep exposed to overpressures in a shock tube. Negaurements of the pH and blood gas tensions, determinations of the venous-admixture (45/4) and the alveolar-arterial oxygen gradient (A-a)C ₂ were conducted before and at intervals up to 132 days following injury. There was an immediate marked increase in Qs/4, reduction in Pa _{O2} , and a moderate increase in (A-a)O ₂ , with very little change in the pH or P _{CO2} of the arterial blood. The greatest recovery was evident within 24 hours with further gradual improvement seem 2, 7, 14, and 21 days after exposure. After the 21st day, most of the animals exhibited virtual complete recovery of the functional efficiency of the pulmonary system as tested at rest.			+	
The pattern of recovery of the respiratory system from blast injury was investigated in sheep exposed to overpressures in a shock tube. Measurements of the pH and blood gas tensions, determinations of the vencus-admixture (w/Q) and the alveolar-arterial oxygen gradient (A-a)C ₂ were conducted before and at intervals up to 132 days following injury. There was an immediate marked increase in Os/Q, reduction in PaO ₂ , and a moderate increase in (A-a)C ₂ , with very little change in the pH or PCO ₂ of the arterial blood. The greatest recovery was evident within 24 hours with further gradual improvement seen 2, 7, 14, and 21 days after exposure. After the 21st day, most of the animals exhibited virtual complete recovery of the functional efficiency of the pulmonary system as tested at rest.				- ·
The pattern of recovery of the respiratory system from blast injury was investigated in sheep exposed to overpressures in a shock tube. Measurements of the pH and blood gas tensions, determinations of the venous-admixture (45/Q) and the alveolar-arterial oxygen gradient (A-a)C ₂ were conducted before and at intervals up to 132 days following injury. There was an immediate marked increase in Qs/Q, reduction in Pa _{O2} , and a moderate increase in (A-a)C ₂ , with very little change in the pH or P _{CO2} of the arterial blood. The greatest recovery was evident within 24 hours with furthe gradual improvement seen 2, 7, 14, and 21 days after exposure. After the 21st day, most of the animals exhibited virtual complete recovery of the functional efficiency of the pulmonary system as tested at rest.		Margurung	u, D.C.	20307
gated in sheep exposed to overpressures in a shock tube. Measurements of the pH and blood gas tensions, determinations of the vencus-admixture (15/2) and the alveolar-arterial oxygen gradient (A-a)C ₂ were conducted before and at intervals up to 132 days following injury. There was an immediate marked increase in 05/2, reduction in Pa _{O2} , and a moderate increase in (A-a)C ₂ , with very little change in the pH or P _{CO2} of the arterial blood. The greatest recovery was evident within 24 hours with furthe gradual improvement seen 2, 7, 14, and 21 days after exposure. After the 21st day, most of the animals exhibited virtual complete recovery of the functional efficiency of the pulmonary system as tested at rest. /			m Auron 101	at indicate sea inserti
blood gas tensions, determinations of the vencus-admixture (45/Q) and the alveolar-arterial oxygen gradient (A-a)C ₂ were conducted before and at intervals up to 132 days following injury. There was an immediate marked increase in Qs/Q, reduction in Pa _{O2} , and a moderate increase in (A-a)C ₂ , with very little change in the pH or P _{CO2} of the arterial blood. The greatest recovery was evident within 24 hours with furthe gradual improvement seen 2, 7, 14, and 21 days after exposure. After the 21st day, most of the animals exhibited virtual complete recovery of the functional efficiency of the pulmonary system as tested at rest.	-			
arterial oxygen gradient (A-a)C ₂ were conducted before and at intervals up to 132 days following injury. There was an immediate marked increase in Qs/Q, reduction in Pa _{O2} , and a moderate increase in (A-a)C ₂ , with very little change in the pH or P _{CO2} of the arterial blood. The greatest recovery was evident within 24 hours with furthe gradual improvement seem 2, 7, 14, and 21 days after exposure. After the 21st day, most of the animals exhibited virtual complete recovery of the functional efficiency of the pulmonary system as tested at rest. /				
days following injury. There was an immediate marked increase in Qs/Q, reduction in PaO2, and a moderate increase in (A-a)O2, with very little change in the pH or POO2 of the arterial blood. The greatest recovery was evident within 24 hours with furthe gradual improvement seen 2, 7, 14, and 21 days after exposure. After the 21st day, most of the animals exhibited virtual complete recovery of the functional efficiency of the pulmonary system as tested at rest.				
Page, and a moderate increase in (A-a)02, with very little change in the pH or PCO2 of the arterial blood. The greatest recovery was evident within 24 hours with furthe gradual improvement seen 2, 7, 14, and 21 days after exposure. After the 21st day, most of the animals exhibited virtual complete recovery of the functional efficiency of the pulmonary system as tested at rest.				
of the arterial blood. The greatest recovery was evident within 24 hours with furthe gradual improvement seen 2, 7, 14, and 21 days after exposure. After the 21st day, most of the animals exhibited virtual complete recovery of the functional efficiency of the pulmonary system as tested at rest.	days following injury. There was a	n immediate marked	increase :	in Qs/Q, reduction in
of the arterial blood. The greatest recovery was evident within 24 hours with furthe gradual improvement seen 2, 7, 14, and 21 days after exposure. After the 21st day, most of the animals exhibited virtual complete recovery of the functional efficiency of the pulmonary system as tested at rest.	Page, and a moderate increase in (A	-a)02, with very li	ttle chang	ge in the pH or PCOo
gradual improvement seen 2, 7, 14, and 21 days after exposure. After the 21st day, most of the animals exhibited virtual complete recovery of the functional efficiency of the pulmonary system as tested at rest. /	75			
most of the animals exhibited virtual complete recovery of the functional efficiency of the pulmonary system as tested at rest. /				
of the pulmonary system as tested at rest. /				
DD FORM 1473			A OI CINE	imiectorist structorich
	or the himonera sastem as rested a	C rest.		
	•			
	<u> </u>	· · · · · · · · · · · · · · · · · · ·		
	DD 104 1473			
Security Classification	WW 1 1997 11 17 / 3	صائبين		

UNCLASSIFIED

Security Classification		والسيساسي			_			_		
14.	KEY WORDS	1.		4 .	100		LIN		LINI	
The same of the sa	** * * * * * * * * * * * * * * * * * *	1			ROLE	WΤ	HOLE	MI	ROLE	WT
atFenent	made solder		;	i .	∤ : ,, ,			$\zeta = \tilde{U}_b$	(I	
Blast Lung Injury	each control				1]	145%			į.
						<u> </u>		.	1 1	
pH, PO ₂ , PCO ₂ in Sheep Venous Admixture							ļ		1 1	
Venous Admixture		5.4.	ř,	:		l		1		
Blood Gas Analysis Pulmonary Function					1	1		ļ .		
Overpressure		****		· V		ŀ	1			
0101220000]	<u>.</u>	1	1		
							1	•		
						1	1	Ì		
						1	1	1)	
					1	ł	!	1		
						1	ļ			
					1	}	Ì	1		
					ļ	ļ	İ	{		
1						1]	
1					1	1				·
4					l	Į.]	ļ	1	
1	,				1				1	
1					1		l "	1	1	,
ŧ						l	İ	1		
					l	1		1		
1					1	}		1	}	
į						1	l	l		
					1	}	l			
1					}	1	1	1		
					1	l	l			
					1	1			<u> </u>	
			•		1	}		1	1	
10					1	1	1	1		
l		•		,	1				j	
1					1	1	1		1	
Į.					1	1	1	1		
1					1		1		1	1
•					1	1	1	1	1	l
· ·		Mr.			1	1		1		1
						1		1	1	·
}					1	1		1		1
I					1	1	1			
1										
1					1	1	1	1	1	!
							1		1	,
1					1	1		1	}	,
1					{	1	1		1	
1					1				1	'
1							1		<u> </u>	

Sounty Closelfinates